



Elbow Locking Caused by Annular Ligament in a Patient with Congenital Radioulnar Synostosis: A Case Report and Literature Review

Konjenital Radioulnar Sinositozlu Hastada Annuler Bağ Hipertrofisine Bağlı Kilitlenme: Olgu Sunumu ve Literatürün Gözden Geçirilmesi

Serkan Aykut¹, Kahraman Öztürk¹, Ayşe Şencan², Fettah Büyük³

Öz / Abstract

Congenital radioulnar synostosis is a rare but challenging congenital elbow anomaly. Patients have severe restriction of forearm rotation, but most patients do not report flexion-extension limitation. Here we report the case of an 11-year-old boy with congenital radioulnar synostosis. He had painful snapping of the right elbow. Painful snapping, and locking occasionally, occurred at the right elbow during flexion of the elbow from extension. There was no history of trauma. Magnetic resonance imaging showed tissue structure between the radial head and capitellum. Open surgery was performed. We found a hypertrophic annular ligament between the radial head and capitellum. During flexion, the ligament got trapped between the radial head and capitellum. The hypertrophic ligament was removed. After the surgery, the patient had no recurrence of snapping or locking.

Keywords: Congenital, radioulnar synostosis, annular ligament

Konjenital Radioulnar sinositoz nadir görülen fakat zorlayıcı konjenital dirsek anomalilerindedir. Hastaların ön kol rotasyonları ciddi olarak kısıtlanmışken çoğu olguda dirsek fleksiyon ve ekstansiyonu etkilenmez. Olgumuz konjenital radioulnar sinositozlu 11 yaşında erkek çocuğuydu. Sağ dirseğinde ani gelişen hareket ile rahatsız edici ses gelmesi şikayeti vardı. Ağrılı sürtünme ve bazen dirseğini ekstansiyondan fleksiyona getirirken kilitlenme oluyordu. Travma öyküsü yoktu. Manyetik rezonans incelemesinde radius başı ve kapitellum arasında yer kaplayan bir doku görüldü. Açık cerrahi uygulandı. Radius başı ve kapitellum arasında annuler bağın hipertrofik olduğu görüldü. Hipertrofik bağ eksize edildi. Cerrahi sonrası hastanın hareket ile sürtünme sesi ve kilitlenme şikayetleri tekrar etmedi.

Anahtar Kelimeler: Konjenital, radioulnar sinositoz, annuler bağ

Introduction

Congenital radioulnar synostosis is a rare disease characterized by a separation anomaly between proximal parts of the radius and ulna (1-3). Frequent family history and association with other congenital syndromes suggest a genetic origin. Proximal parts of the radius and ulna are typically fused in these patients (1). Forearm rotation is restricted in these patients, but elbow motion is generally preserved except for patients with flexion contracture (4, 5).

In this study, we aimed to present a patient with congenital radioulnar synostosis, a rare condition that resulted in recurrent elbow locking and crepitation on range of motion.

Case Report

An 11-year-old, right-hand dominant boy was brought to our emergency department with the complaint of a painful locked elbow. His past medical history included emergency admission three times due to elbow locking. Despite medical treatment and activity restriction, patient's complaints did not regress. He had right congenital proximal radioulnar synostosis. He had neither trauma nor any history of operation of the elbow. Inspection did not reveal any ecchymosis, edema, or rash. Active extension and flexion were limited to -20° and 110°, respectively. The right forearm was in a fixed pronated position at 10°. Right elbow flexion was painful with crepitation during range of motion. Additionally, there was a history of locking in during sudden elbow movements. Locking was not elicited during examination. Flexion in the left elbow ranged between 0° and 140°; the forearm was in a fixed pronated position at 10°. Radiographs revealed an anterior radial head dislocation and a type 4 congenital radioulnar synostosis according to Cleary and Omer classification (1). Magnetic resonance imaging (MRI) showed a hypertrophic space-occupying lesion between the radius head and capitellum. MR arthrogram was not performed. Debridement of the lesion, which was the cause of the locking, with an open surgery was recommended. The patient underwent open surgery under general anesthesia. Under tourniquet control, lateral approach to the elbow was performed. By partial detachment of the extensor muscle origin and releasing the anterior joint capsule from the humerus, the radiohumeral joint was exposed. When the elbow was moved into the flexion from extension position, a fibrous band thought to be a remnant of the annular ligament, caused crepi-

¹Department of Hand and Upper Extremity Surgery, Metin Sabancı Baltalimanı Bone Diseases Training and Research Hospital, Istanbul, Turkey

²Department of Hand Surgery, Bağcılar Training and Research Hospital, Istanbul, Turkey

³Department of Orthopedics and Traumatology, Haseki Training and Research Hospital, Istanbul, Turkey

Address for Correspondence

Yazışma Adresi:

Serkan Aykut

E-mail: aykutserkan@yahoo.com

Received/Geliş Tarihi: 28.03.2016

Accepted/Kabul Tarihi: 26.04.2017

© Copyright 2017 by Available online at www.istanbulmedicaljournal.org

© Telif Hakkı 2017 Makale metnine www.istanbultipdergisi.org web sayfasından ulaşılabilir.



Figure 1. During surgery, the hypertrophic annular ligament can be seen when the elbow is in the flexion position from extension. The annular ligament was causing crepitation and elbow locking by getting trapped between the radial head and capitellum



Figure 2. Hypertrophic annular ligament was excised with preservation of the lateral collateral ligament

tation and locking by getting trapped between the radial head and capitellum (Figure 1). The hypertrophic tissue was excised with the preservation of the lateral collateral ligament (Figure 2). Crepitation resolved after the excision. Range of motion exercises were immediately initiated after surgery without cast immobilization. Histological evaluation showed ligament structure and no evident synovial tissue. Three years after the surgery, the patient has no complaints and no recurrence of elbow locking. Written informed consent was received from the parents of the patient for the publication of the case report.

Discussion

Acute elbow locking is often due to joint mouse. However, it may also be seen due to extra-articular causes, such as annular ligament subluxation, ulnar nerve dislocation, and snapping of medial head of the triceps, or in intra-articular causes, such as synovial plica impingement and posterolateral rotator instability (6-9). MRI has limited diagnostic value in patients with crepitation or elbow locking. MR arthrography is the preferred imaging modality in identifying anatomical abnormalities inside the joint (5, 7). The most valuable diagnostic and therapeutic tool is the arthroscopic approach. This approach allows the intra-articular pathology to be directly observed, and elbow dynamic examination to be evaluated. Pathologic tissues can be removed with this tool (5, 6, 10).

This is the third case report of elbow locking in a patient with radioulnar synostosis in the English literature. Masuko et al. (3) reported two radioulnar synostosis cases with elbow flexion contracture due to the presence of annular ligament-like tissue. They described good outcome with excision of the annular ligament-like tissue, as we performed in our case. Shinora et al. (5) operated on two patients: one presented with painful snapping and the other with elbow locking. They found a tight fibrous tissue trapping the radial head, and arthroscopic removal of the fibrous tissue resolved the symptoms. Shin et al. (4) reported one radioulnar synostosis case with limitation of elbow flexion due to enlarged radial head. They excised approximately 15 mm of the radial head and 6 months later, there was full flexion of the elbow.

Conclusion

Annular ligament-like fibrous band is rarely seen in patients with congenital radioulnar synostosis, but it can result in elbow locking, and should, therefore, be considered for elbow locking.

Informed Consent: Written informed consent was obtained from the patient who participated in this study.

Peer-review: Externally peer-reviewed.

Author contributions: Concept – S.A., K.Ö.; Design – S.A., F.B., A.Ş.; Supervision – S.A., K.Ö.; Resource - S.A.,F.B.; Materials - S.A., F.B.; Data Collection and/or Processing - S.A., F.B.; Analysis and/or Interpretation - S.A., K.Ö.; Literature Search - S.A., F.B.; Writing - S.A., F.B.; Critical Reviews - S.A., F.B., A.Ş., K.Ö.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Hasta Onamı: Yazılı hasta onamı bu çalışmaya katılan hastadan alınmıştır.

Hakem Değerlendirmesi: Dış Bağımsız.

Yazar Katkıları: Fikir – S.A., K.Ö.; Tasarım - S.A., F.B., A.Ş.; Denetleme - S.A.,K.Ö.; Kaynaklar - S.A.,F.B.; Malzemeler - S.A., F.B.; Veri Toplanması ve/veya İşlemesi - S.A., F.B.; Analiz ve/veya Yorum - S.A., K.Ö.; Literatür taraması - S.A., F.B.; Yazıyı Yazan - S.A., F.B.; Eleştirel İnceleme - S.A., F.B., A.Ş., K.Ö.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

References

1. Cleary JE, Omer GE. Congenital proximal radio-ulnar synostosis natural history and functional assessment. *J Bone Joint Surg* 1985; 67: 539-45. [\[CrossRef\]](#)
2. Ramachandran M, Lau K, Jones DHA. Rotational osteotomies for congenital radioulnar synostosis. *J Bone Joint Surg Br* 2005; 87: 1406-10. [\[CrossRef\]](#)
3. Masuko T, Kato H, Minami A, Inoue M, Hirayama T. Surgical Treatment of acute elbow flexion contracture in patients with congenital proximal radioulnar synostosis a report of two cases. *J Bone Joint Surg Am* 2004; 86: 1528-33. [\[CrossRef\]](#)
4. Shin YH, Baek GH, Lee HJ. Limitation of elbow flexion in a patient with congenital radioulnar synostosis. *J Hand Surg* 2012; 37: 576-7. [\[CrossRef\]](#)

5. Shinohara T, Horii E, Tatabe M, Yamamoto M, Okui N, Hirata H. Painful snapping elbow in patients with congenital radioulnar synostosis: report of two cases. *J Hand Surg* 2010; 35: 1336-9. [\[CrossRef\]](#)
6. Steinert AF, Goebel S, Rucker A, Barthel T. Snapping elbow caused by hypertrophic synovial plica in the radiohumeral joint: a report of three cases and review of literature. *Acta Orthop Trauma Surg* 2010; 130: 347-51. [\[CrossRef\]](#)
7. Huang GS, Lee CH, Lee HS, Chen CY. A meniscus causing painful snapping of the elbow joint: MR imaging with arthroscopic and histologic correlation. *Eur Radiol* 2005; 15: 2411-4. [\[CrossRef\]](#)
8. O'Driscoll SW. Classification and evaluation of recurrent instability of the elbow. *Clin Orthop Relat Res* 2000; 370: 34-43. [\[CrossRef\]](#)
9. Watts AC, McEachan J, Reid J, Rymaszewski L. The snapping elbow: A diagnostic pitfall. *J Shoulder Elbow Surg* 2009; 18: 9-10. [\[CrossRef\]](#)
10. Maruyama M, Takahara M, Kikuchi N, Ito K, Watanabe T, Ogino T. Snapping elbow with congenital radial head dislocation: case report. *J Hand Surg* 2010; 35: 981-5. [\[CrossRef\]](#)

Cite this article as: Aykut S, Öztürk K, Şencan A, Büyük F. Elbow Locking Caused by Annular Ligament in a Patient with Congenital Radioulnar Synostosis: A Case Report and Literature Review. *Istanbul Med J* 2017; 18: 242-4.