



A Case of Sporadic Creutzfeldt-Jakob Disease Mimicking Variant Form

Varyant Formu Taklit Eden Sporadik Creutzfeldt-Jakob Hastalığı Olgusu

Osman Özgür Yalın, İrem Baş, Ufuk Emre, Orhan Yağız, Yüksel Erdal

Öz / Abstract

Creutzfeldt-Jakob Disease (CJD) is a rare neurodegenerative disease that causes a rapidly progressive dementia, and it can be classified as sporadic, heritable, and variant (acquired). In this paper, we report a case mimicking variant CJD, with phenotypic presentation and findings. A 44-year-old man was admitted to the emergency department with progressive neuropsychiatric symptoms and myoclonus. Signal hyperintensity in the bilateral basal ganglia and thalamus was detected via fluid attenuated inversion recovery (FLAIR) sequence and diffusion weighted magnetic resonance. 14.3.3 protein in cerebrospinal fluid was negative, and electroencephalography was atypical for sporadic CJD. After the diagnostic tests and follow-up, the patient was diagnosed with probable sporadic CJD, based on the World Health Organization's (WHO) diagnostic criteria. His health deteriorated rapidly, and the patient died 2 months after the diagnosis. In this paper, we point out the differences between and the signs of variant CJD and sporadic CJD. When pathological confirmation is not possible, as in case of our patient, it is very helpful to consider the WHO diagnostic criteria. Variant CJD is a disorder that has not been reported in Turkey yet. Its diagnosis and reporting require direct interventions by the health and veterinary authorities.

Keywords: Prion, variant Creutzfeldt-Jakob Disease, pulvinar sign

Creutzfeldt-Jacob Hastalığı (CJH) hızlı progresif demansa neden olan, sporadik, kalıtsal ve varyant (edinilmiş) formları olan nadir görülen bir nörodejeneratif hastalıktır. Bu yazıda fenotipik prezentasyon ve muayene bulgularıyla varyant CJH'yi taklit eden bir olgu sunduk. 44 yaşında erkek hasta, ilerleyici nöropsikiyatrik şikayetler ve miyoklonus bulguları ile acil servisimize başvurdu. FLAIR ve difüzyon ağırlıklı manyetik rezonans görüntülemelerde, bilateral bazal ganglion ve talamusta sinyal artışı saptandı. Beyin omurilik sıvısında 14.3.3 protein negatif ve elektroensefalografi sporadik CJH için tipikti. Tanıya yönelik incelemeler ve takipleri sonrasında, dünya sağlık örgütü (WHO) tanı kriterlerine dayanılarak muhtemel sporadik CJH tanısı aldı. Hasta hızla kötüleşti ve tanı konulduktan iki ay sonra hayatını kaybetti. Bu yazımızda varyant CJH ve sporadik CJH'nin farklılıklarına ve tanılarda ipuçlarına dikkat çektiğimiz hastamız gibi patolojik tanıya erişilemediğinde DSÖ tanı ölçütlerini hatırlamak çok faydalı olacaktır. Varyant CJH, Türkiye'de henüz rapor edilmediği vurgulanan bir hastalıktır, teşhis ve raporlamak için hükümetin sağlık ve veterinerlik makamlarının doğrudan müdahaleleri gerekir.

Anahtar Kelimeler: Prion, varyant Creutzfeldt-Jacob Hastalığı, pulvinar sign

Introduction

Creutzfeldt-Jacob disease (CJD) is a rare, rapid, and progressive fatal human prion disease, affecting annually 1-2 persons per million worldwide. In addition to abnormal protein accumulation, CJD is characterized by neuronal loss, gliosis, and spongiform changes. CJD is classified as sporadic, genetic, iatrogenic, and variant CJD (1). In this report, we will discuss a patient on phenotypic expression of sporadic and variant CJD.

Whereas the majority of CJD cases (about 85%) occur as sporadic disease, clinical presentation of sporadic CJD is highly variable. CJD shows a broad range of symptoms, but the disease begins with cognitive and cerebellar symptoms in most of cases. Visual changes and extrapyramidal and pyramidal signs are less frequently seen as initial manifestations (1, 2). Myoclonus occurs during the clinical course in approximately 90% of cases, and the disease eventually progresses to akinetic mutism in all patients. In most of cases, the disease lasts less than a year, but the course of sporadic CJD depends on polymorphisms in PRNP codon 129 and the type of protease-K resistant prion (3).

Variant CJD was identified in 1996 attracting attention as a casual link to bovine spongiform encephalopathy (4). Variant CJD is still the only example of an infectious disease crossing from bovines to humans. In contrast to the sporadic CJD, clinical presentation of variant CJD is unique almost in all cases, and diagnosis could be predicted confidently by clinical findings and neuroimaging (5). Persistent sensory symptoms, early psychiatric symptoms, chorea, and upgaze paresis are much more common in variant CJD, whereas early forgetfulness and ataxia are relatively uncommon (6).

Istanbul Eğitim ve Araştırma Hastanesi, Nöroloji Kliniği, İstanbul, Türkiye

Address for Correspondence


Yazışma Adresi:
Osman Özgür Yalın
E-mail: osmanozguryalin@yahoo.com

Received/Geliş Tarihi: 25.01.2017

Accepted/Kabul Tarihi: 07.05.2017

© Copyright 2017 by Available online at
www.istanbulmedicaljournal.org

© Telif Hakkı 2017 Makale metnine
www.istanbultipdergisi.org web sayfasından
ulaşılabilir.



Resim 1. a-c. Brain magnetic resonance imaging (MRI) shows bilateral striatal and thalamic hyperintensities. (a) Fluid attenuated inversion recovery (FLAIR) sequence and, (b) diffusion weighted imaging show bilateral striatal and thalamic hyperintensities, (c) apparent diffusion coefficient map shows striatal and mildly thalamic hypointensity. Also note that anterior putamen and caudate are more hyperintense than the thalamus

Case Report

A 44-year-old man was admitted to our emergency department complaining of dizziness, fatigue, and difficulties at walking. The history revealed complaints that started 2 months before. Dizziness and behavioral symptoms started 2 weeks after the upper-respiratory-tract infection. There were occult behavioral symptoms present, including agitation, anxiety, and bizarre new ideas, and there was not past psychiatric disorder history. Patient had been using ketiapin 200 mg/day for a week, as recommended by a psychiatric specialist, and he was admitted to our hospital due to worsening of symptoms, as proposed by a psychiatrist. There was no past illness history and family history of dementia or a neuropsychiatric disease. During the neurologic examination, cooperation and orientation were normal, marked photophobia was present, whereas cranial nerve examination showed bilateral upper-gaze palsy and prominent truncal ataxia. Myoclonic jerks were present in the upper extremities and facial muscles, particularly periocular muscles, nearly continuous. The patient was hospitalized, and magnetic resonance imaging (MRI) and laboratory tests were performed. MRI diffusion weighted and fluid attenuated inversion recovery (FLAIR) sequence showed hyperintensity at symmetrically bilateral thalamus, and caudate and lentiform nucleus (Figure 1). Laboratory findings, including serum biochemistry, hemogram, liver and kidney function test, thyroid function test, vitamin B12 and folate levels, were within the normal limits. Electroencephalography (EEG) showed alpha and theta rhythm and generalize slowing, and there was no periodic sharp continuous wave activity and/or other epileptiform discharge. Lumbar puncture was performed; cerebrospinal fluid (CSF) opening pressure, glucose, and protein levels were within a normal range, and 14.3.3 protein was negative. Patient's condition deteriorated rapidly, and at the 2-month follow-up, akinetic mutism developed. The patient was fed through percutaneous endoscopic gastrostomy catheter. Two weeks later, the patient was transferred to the intensive care unit because of a respiratory failure, and he died 2 months later. Written consent was obtained from patient's relatives.

Discussion

Our goal was to discuss this case in the light of phenotypic characteristics and features that distinguish sporadic CJD from variant CJD. The young age at onset, early presentation with psychiatric disturbances, upgaze paresis, atypical EEG, negative 14.3.3 protein in CSF and MRI diffusion hyperintensities at thalamus, and caudate and lentiform nucleus alerted us for variant CJD probability. To our knowledge, there is only one probable case of variant CJD in Turkey reported in 2011 without pathological confirmation (7). We did not consider the MRI of our patient to be concordant with variant CJD. The WHO listed diagnostic criteria for variant CJD in 2001 (6). In this report, definite variant CJD diagnosis is based on a progressive neuropsychiatric disorder and pathologic confirmation. Variant CJD diagnosis is probable if (a) the illness lasts longer than 6 months and other typical clinical signs for variant CJD occur; (b) EEG does not show the clinical characteristics of sporadic CJD; and (c) MRI shows bilateral symmetrical pulvinar high signals. The most important segment of the MRI criteria is "pulvinar sign." Although other conditions may also cause thalamic and other gray matter hyperintensities, in variant CJD, the highest signal is always in the pulvinar of the thalamus, and this is an essential distinguishing feature. The pulvinar sign has a 90% sensitivity for the diagnosis of variant CJD, and the specificity of the pulvinar sign is over 95%. The pulvinar sign is the best non-invasive in vivo diagnostic test for variant CJD. Will R. G. reported that all the cases fulfilling these criteria, including MRI features for probable variant CJD, have been confirmed pathologically as variant CJD (4). Although there is one case of probable variant CJD reported in Turkey (7), the MRI did not show the characteristic "pulvinar sign." Will RG (8) also discussed this report and assumed that this case cannot be regarded as variant CJD unless there is a neuropathological verification. Thus far, Turkey is geographically regarded as variant-CJD unreported area.

With regard to our case, we assumed that it was sporadic CJD with some peculiar findings. These findings could be attributed to young-age onset or a molecular subtype based on the polymorphisms in PRNP codon 129 (2). We could not perform autopsy because the family did not approve it. We did not plan tonsil biopsy because neu-

ropsychiatric complaints lasted less than 6 months, although other clinical findings, EEG and CSF, were atypical for sporadic CJD.

Conclusion

It is important to diagnose and report variant CJD in our country if it exists, because even one case of this infectious lethal disease would require a prompt action by state health and veterinary authorities. Ultimately, we believe and point out that when it is not possible to perform biopsy, like in case of our patient, the WHO diagnostic criteria could prevent misdiagnosis of variant CJD. Diagnostic criteria are especially important in countries where performing an autopsy is challenging.

Informed Consent: Written informed consent was obtained from the relatives of the patient who participated in this study.

Peer-review: Externally peer-reviewed.

Author contributions: Concept - O.O.Y., İ.B., U.E.; Design - O.O.Y., İ.B., U.E.; Supervision - U.E., O.Y.; Resource - O.O.Y., İ.B., U.E.; Materials - O.O.Y., İ.B.; Data Collection and/or Processing - İ.B., O.O.Y., Y.E.; Analysis and/or Interpretation - O.Ö.Y., U.E., Y.E.; Literature Search - O.Ö.Y., İ.B., U.E.; Writing - O.O.Y., İ.B., Y.E.; Critical Reviews - U.E., O.Y.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Hasta Onamı: Yazılı hasta onamı çalışmaya katılan hastanın yakınlarından alınmıştır.

Hakem Değerlendirmesi: Dış Bağımsız.

Yazar Katkıları: Fikir - O.O.Y., İ.B., U.E.; Tasarım - O.O.Y., İ.B., U.E.; Denetleme - U.E., O.Y.; Kaynaklar - O.O.Y., İ.B., U.E.; Malzemeler - O.O.Y., İ.B.;

Veri Toplanması ve/veya işlemesi - İ.B., O.O.Y., Y.E.; Analiz ve/veya Yorum - O.Ö.Y., U.E., Y.E.; Literatür taraması - O.Ö.Y., İ.B., U.E.; Yazıyı Yazan - O.O.Y., İ.B., Y.E.; Eleştirel İnceleme - U.E., O.Y.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemiştir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

References

1. Takada LT, Geschwind MD. Prion diseases. *Semin Neurol* 2013; 33: 348-56. [\[CrossRef\]](#)
2. Corato M, Cereda C, Cova E, Ferrarese C, Ceroni M. Young-onset CJD: age and disease phe-notype in variant and sporadic forms. *Funct Neurol* 2006; 21: 211-5.
3. Puoti G, Bizzi A, Forloni G, Safar JG, Tagliavini F, Gambetti P. Sporadic human prion diseases: molecular insights and diagnosis. *Lancet Neurol* 2012; 11: 618-28. [\[CrossRef\]](#)
4. Will RG, Ironside JW, Zeidler M, Cousens SN, Estibeiro K, Alperovitch A, et al. A new variant of Creutzfeldt-Jakob disease in the UK. *Lancet* 1996; 347: 921-5. [\[CrossRef\]](#)
5. Will RG, Zeidler M, Stewart GE, Macleod MA, Ironside JW, et al. Diagnosis of New Variant Creutzfeldt-Jakob Disease. *Ann Neurol* 2000; 47: 575-82. [\[CrossRef\]](#)
6. World Health Organization: The Revision of the Surveillance Case Definition for Variant Creutzfeldt-Jakob Disease (vCJD). Report of a WHO consultation Edinburgh, United Kingdom. 2001: 5.
7. Adanipar DÖ, Saylısoy S, Yenilmez Ç, Aslan H, Ertan B, Artan S, et al. The first report of a patient with probable variant Creutzfeldt-Jakob disease in Turkey. *Dement Geriatr Cogn Dis Extra* 2011; 1: 429-32. [\[CrossRef\]](#)
8. Will RG. Response to the Article by Adanipar et al.: The First Report of a Patient with Pro-bable Variant Creutzfeldt-Jakob Disease in Turkey. *Dement Geriatr Cogn Disord Extra* 2012; 2: 81-2.

Cite this article as: Yalın OÖ, Baş İ, Emre U, Yağız O, Erdal Y. A Case of Sporadic Creutzfeldt-Jakob Disease Mimicking Variant Form. *İstanbul Med J* 2017; 18: 245-7.