



Short-term Quality of Life after Cytoreductive Surgery and Hyperthermic Intraperitoneal Chemotherapy

Sitoredüktif Cerrahi ile İntraperitoneal Sıcak Kemoterapinin Erken Dönem Yaşam Kalitesi Üzerine Etkisi

Özgül Düzgün¹, İnanç Şamil Sarıcı², Serkan Gökçay³

Abstract / Öz

Introduction: Cytoreductive surgery (CRS) and hyperthermic intraperitoneal chemotherapy (HIPEC) have been increasingly performed in many advanced centers of our country. Short-term quality of life (QoL) after the CRS with HIPEC was investigated in Turkish population. The purpose of our study was to investigate the QoL in patients operated for peritoneal carcinomatosis (PC).

Methods: Retrospectively collected data from 42 consecutive patients, who had undergone CRS and HIPEC, and 92 abdominal malignancy patients after oncological surgery without CRS and HIPEC were selected in the study between 2012 and 2015. A Turkish version of the European Organization for Research and Treatment of Cancer questionnaire (QLQ-C30) was used at the 6th month following surgery for the QoL assessment. Statistical analysis was done with one sample t-test, and $p < 0.05$ was accepted as statistically significant.

Results: In total, 42 CRS+HIPEC patients were analyzed; of which, 30 (71.4%) were female and 12 (28.5%) were male. The median age was 52 years (range, 22-69). CRS with HIPEC was performed for colorectal carcinoma in 15 patients (35.7%), ovarian cancer in 12 patients (28.5%), gastric cancer in 6 patients (14.2%), pseudomixoma peritonei in 3 patients (7.1%), sarcoma in 3 patients (7.1%), and mesenchymal tumor in 3 patients (7.1%). The median intraoperative peritoneal carcinomatosis index score was 15 (6-29), whereas the completeness of the cytoreductive score was 0 and 1. The median follow-up period for all patients was 12 months (range, 6-22 months). The functional and symptom scores were equal between the CRS with HIPEC and without CRS+HIPEC cancer patients ($p > 0.05$).

Conclusion: Short-term QoL after the CRS with HIPEC patients and without the CRS+HIPEC oncologic patients is found to be similar in Turkish population.

Keywords: Cytoreductive surgery (CRS), hyperthermic intraperitoneal chemotherapy (HIPEC), Turkish population, quality of life (QoL)

Amaç: Sitoredüktif cerrahi (CRS) ile birlikte hipertermik intraperitoneal sıcak kemoterapi (HIPEC) ülkemizde ileri merkezlerde son yıllarda giderek artan sıklıkla yapılmaktadır. Çalışmamızın amacı peritoneal karsinomatozis (PK) teşhisiyle ameliyat edilen hastaların hayat kalitesi (QoL) değerlendirmektir.

Yöntemler: Çalışmamızda 2012-2015 yılları arasında PK tanısıyla SRC+HIPEC yapılan 42 hastanın ameliyat sonrası hayat kaliteleri ile SRC+HIPEC olmaksızın onkolojik cerrahi sonrası takip edilen 92 hastanın hayat kaliteleri karşılaştırıldı. Avrupa Kanser Araştırma ve Tedavi Organizasyonu (EORTC) tarafından hazırlanan QLQ-C30 soruları tüm hastalarımıza ameliyat sonrası 6. ayda uygulandı ve her iki grup karşılaştırıldı. İstatistiksel analiz one sample t-test ile yapılarak, $p < 0.05$ anlamlı kabul edildi.

Bulgular: 42 hastanın 30'u (%71.4) bayan, 12'si (%28.5) erkek idi. Hastaların yaş ortalaması 52 (range, 69-22) idi. PK orijinleri 15 hastada (%35.7) kolorektal malignite, 12 hastada (%28.5) over kanseri, 6 hastada (%14.2) mide malignitesi, 3 hastada (%7.1) pseudomixoma peritonei, 3 hastada (%7.1) sarkom ve 3 hastada (%7.1) da mezenkimal tümör olarak bulundu. Ortalama intraoperatif peritoneal karsinomatoz indeksi (PCI) 15 (range 6-29) olarak bulundu. Rezeksiyon tamlik skoru (CCS) 0 ve 1 olarak bulundu. Tüm hastalarda ortalama takip süresi 12 ay (range 2-26) idi. Her iki grup arasında fonksiyonel ve semptom skorları açısından istatistiksel fark saptanmadı ($p > 0.05$).

Sonuç: SRC+HIPEC olmaksızın onkolojik cerrahi yapılan hastalar ile karşılaştırıldığında, PK sebebiyle SRC+HIPEC yapılan olguların kısa dönemde yaşam kalitelerinin (QoL) benzer olduğu bulunmuştur.

Anahtar Kelimeler: Sitoredüktif cerrahi (CRS), Hipertermik intraperitoneal kemoterapi (HIPEC), Turkish population, Yaşam kalitesi (QoL)

This study was presented 20th National Surgery Congress (April 13-17, 2016, Antalya, Türkiye)

¹Department of Oncological Surgery, Çukurova University School of Medicine, Adana, Türkiye

²Department of General Surgery, Kanuni Sultan Süleyman Training and Research Hospital, İstanbul, Türkiye

³Department of Medical Oncology, Çukurova Üniversitesi School of Medicine, Adana, Türkiye

Address for Correspondence/Yazışma Adresi:
İnanç Şamil Sarıcı
E-mail: issarici2015@gmail.com

Received/Geliş Tarihi: 28.03.2017

Accepted/Kabul Tarihi: 23.11.2017

© Copyright 2018 by Available online at istanbulmedicaljournal.org

© Telif Hakkı 2018 Makale metnine istanbulmedicaljournal.org web sayfasından ulaşılabilir.

Introduction

Peritoneal carcinomatosis (PC) which is called advanced stage of abdominal cancer continues to be a problem that not fully solved at the present time. Due to both medical cost burdens and inferior treatment results of oncologic diseases with palliative chemo-radiotherapy or supportive care in advance, in 1989 Sugarbaker et al. (1) presented cytoreductive surgery (CRS) and hyperthermic intraperitoneal chemotherapy (HIPEC) as a stunning method for appropriate patients with PC. Initially, this technique had excessive treatment-related morbidity and mortality with a major concern. However, completion of learning curve in the experienced centers resulted an overall decrease in postoperative morbidity and mortality, as shown in the recent studies (2, 3).

Regardless of enhanced survival with admissible morbidity and mortality of surgery, quality of life (QoL) in patients with PC is still controversial. Due to broad resection and possible chemotherapy toxicity, changes in QoL patients with CRS and HIPEC could be more than simple surgery.

These studies were performed in Western and Asian population; nevertheless, have not been applied in Turkish population. Thus, in the present study, we aimed to share out QoL results of our

PC patients with CRS and HIPEC with comparison the oncologic patients operated only without CRS and HIPEC.

Methods

Prospectively collected data of 42 PC patients who had undergone CRS and HIPEC in Department of Surgical Oncology, Çukurova University School of Medicine, between

December 2012 and May 2015, were analyzed. Preoperative evaluation was made with thoraco-abdominal computerized tomography (CT) and positron emission tomography (PET)-CT. Patients were chosen as candidates for CRS and HIPEC in multi-disciplinary oncology meeting aiming a complete cytoreduction. For comparison, 92 abdominal malignancy patients after oncological surgery without CRS and HIPEC were selected randomly in our oncology clinic. Ethic Committee approval was approved by the Institutional Review Board of the Cukurova University for this study (CUM 2015-48-28). Informed consent was obtained from each patient.

Intraoperatively spreading of tumour was evaluated by the PC index (PCI) (4). Residual tumor presence after cytoreduction was reevaluated with residual completeness scoring (CCS) (5). Mitomycin C and cisplatin were used according to the primary origin of the tumor. Chemotherapeutic drug selection and preparation were done by the clinical oncology experts. After CRS, four intraperitoneal drainage catheters were inserted into the abdomen, and HIPEC was administered for 60 minutes (range, 30-60).

The Turkish form of the European Organization for Research and Treatment of Cancer (EORTC) QLQ-C30 which was confirmed previously by Hoopman et al. (6) was selected for QoL. All questionnaires were administered by clinical oncologists. Functional measurement (physical, role, cognitive, emotional, social), symptom degree (fatigue, pain, dyspnea, loss of appetite, sleeping, diarrhea, constipation, nausea or vomiting, and financial problems), and global health status were measured in all patients. Scoring was ranged between 0 and 100 (7).

Statistical Analysis

Statistical analyses were performed using SPSS (Statistical Package for Social Sciences) 15.0 for Windows (SPSS Inc.; Chicago, IL, USA). The mean QoL scores were calculated in both groups and compared by using one-sample t-test. P values <0.05 were considered statistically significant.

Results

Forty-two patients with CRS and HIPEC with 92 oncologic patients operated without CRS and HIPEC were compared. Of the 42 patients with PC, 30 patients (71.4%) were female and 12 (28.5%) were male. The median age was 52 years (range, 22-69). No significant difference was observed according to the demographics of the two groups (Table 1).

CRS with HIPEC were performed for colorectal carcinoma in 15 patients (35.7%), ovarian cancer in 12 patients (28.5%), gastric cancer in 6 patients (14.2%), pseudomyxoma peritonei in 3 patients (7.1%), sarcoma in 3 patients (7.1%), and mesenchymal tumor in 3 patients (7.1%). The median intraoperative peritoneal carcinomatosis index (PCI) score was 15 (6-29) while the completeness of cytoreductive score (CCS) was 0 and 1. The median follow-up pe-

Table 1. Demographics of patients, comorbidity and the primary tumor

Variables	CRS+HIPEC patients N=42	Oncology patients operated without CRS+HIPEC N=92
Gender (M/F)	42(12/30)	92 (31/61)
Age (year)	52 (22-69)	58(25-72)
BMI (kg/m ²)	21.1±4.2	22.2±3.5
Education N (%)		
Primary	4 (9.5 %)	8 (8.7 %)
High School	12 (28.5%)	27 (29.3 %)
University	26 (62 %)	52 (56.5 %)
Comorbidities N		
Hypertension	12	14
Diabetes mellitus	7	17
Asthma	3	5
Cardiovascular disease	1	6
Primary tumor N (%)		
Colorectal	15(35.7%)	44(47.8%)
Ovarian	12(28.5%)	30(32.6%)
Gastric	6(14.2%)	15 (16.3 %)
Pseudomyxoma peritonei	3 (7.2 %)	-
Sarcoma	3 (7.2 %)	3 (3.2 %)
Mesenchymal tumor	3 (7.2 %)	-

*CRS: cytoreductive surgery; HIPEC: hyperthermic intraperitoneal chemotherapy; BMI: body mass index

riod for all patients was 12 months (range, 6 - 22). Functional and symptom scores were equal between CRS with HIPEC and without CRS+HIPEC cancer patients ($p>0.05$). The ECOG performance status of all patients was 0 or 1.

Without PC group included colorectal carcinoma in 44 patients (47.8%), ovarian cancer in 30 patients (32.6%), gastric carcinoma in 15 patients (16.3%), and sarcomas in 3 patients (3.2%).

The median duration of operation for CRS and HIPEC was 480 minutes (range, 310-565) and that for without CRS and HIPEC group was 110 minutes (range, 80-160) ($p<0.001$). The median PCI was 15 (range, 6-29). There were 39 patients in CCS 0 and 3 patients in CCS 1. The median admission in intensive care unit (ICU) was 1 day (range, 0-12). Median length of hospital stay was 13 days (range, 7-28).

The EORTC-QLQ-C30 scores of patients after CRS and HIPEC compared with patients without CRS and HIPEC are demonstrated in Table 2. There were no difference between the groups of EORTC-QLQ-C30 scores.

Discussion

CRS and HIPEC enhanced survival of patients with peritoneal dissemination of cancer (1-3). Despite improved survival and admissible morbidity and mortality of the surgery, QoL for the patients undergoing CRS and HIPEC is still controversial. In spite of there

Table 2. Comparison of EORTC QLQ C-30 scores in patients with CRS and HIPEC and oncology patients operated without CRS and HIPEC

EORTC QLQ-C30 parameters	CRS+HIPEC patients (N=42)	Oncology patients operated without CRS+HIPEC (N=92)	P
Physical functioning	78(70-87)	84 (75-90)	0.524
Role functioning	81(76-91)	85 (79-93)	0.406
Cognitive functioning	87(81-94)	89 (85-96)	0.305
Emotional functioning	82(75-88)	84 (73-90)	0.555
Social functioning	81(75-90)	82 (75-90)	0.465
Global quality of life	65(57-79)	68 (55-82)	0.343
Fatigue	14(10-22)	16 (11-25)	0.245
Pain	15(6-24)	16 (7-25)	0.324
Nausea and vomiting	8(2-15)	9 (1-17)	0.405
Dyspnea	10(3-17)	9 (2-17)	0.325
Insomnia	19(8-28)	17 (8-26)	0.308
Appetite loss	10(2-18)	9 (3-17)	0.243
Constipation	15(6-22)	13 (5-21)	0.543
Diarrhea	9 (3-15)	8(2-17)	0.550
Financial problems	23(11-35)	24 (11-37)	0.605

*CRS: cytoreductive surgery; HIPEC: hyperthermic intraperitoneal chemotherapy; EORTC QLQ-C30: European Organization for research and treatment of cancer quality of life

have been many articles with respect to QoL scoring calculated for patients with CRS and HIPEC, there is no information about these scores in Turkish PC patients. To our knowledge, this is the first study from a Turkish university hospital referral oncology unit that assesses QoL outcomes who undergone CRS with HIPEC.

There have been recent publications in the literature about EORTC QLQ C-30 to assess the QoL for PC patients. Schmidt et al. (8) evaluated the QoL of patients who underwent CRS combined with HIPEC, and found that QoL is associated with an increased morbidity and mortality according to the difficulty of surgery. On the other hand, in the study of Alves et al. (9) applied EORTC QLQ C-30 questionnaire to their PC patients who undergone CRS with HIPEC. The authors found an improvement in QoL at one year following the procedure. Furthermore, Tan et al. (10) compared CRS plus HIPEC patients with their cancer free patients. They found cognitive functioning scores and fatigue scores better in their CRS+HIPEC patients.

In the present study, the Turkish version of EORTC QLQ C-30 was evaluated between PC patients with CRS and HIPEC and patients without CRS and HIPEC, 6 months after surgery, and no significant difference was found between two groups. However, our study have some limitations. First one is its retrospective design. Lack of the patients basic QoL scores for reference values after CRS and HIPEC is the second limitation. Third limitation is the variety of primary cancer origins. Compared to the population of oncology patients operated without CRS and HIPEC in our center, those patients that underwent CRS and HIPEC had similar scores in global health, physical functioning, emotional functioning and social functioning.

Developing hospital care, advances in ICU for postoperative patient care, and a significant increase in the numbers of specialized oncology units in private sector as well as in the state hospitals can

be the reason of no difference of QoL. On the other hand, successful tumor free surgery, lack of serious morbidity, adequate enteral nutrition, achievement of longer disease-free survival and rapid return to normal life of the oncology patients are important factors affecting QoL. This is likely due to the fact that many of the patients after cancer surgery in Turkey starting to take chemotherapy in this period, quick return to their normal economic, social and physical lives that can be explained why scores are better in both CRS and HIPEC and without CRS and HIPEC groups.

Conclusion

Short term QoL after CRS with HIPEC and oncology patients operated without CRS and HIPEC seems to be equal in our Turkish population.

Ethics Committee Approval: Ethic Committee approval was received by the Institutional Review Board of the Çukurova University for this study (CUM 2015-48-28).

Informed Consent: Informed consent was obtained from the patients who participated in this study

Peer-review: Externally peer-reviewed.

Author contributions: Concept - Ö.D. İ.Ş.S.; Design - İ.Ş.S, Ö.D., S.G.; Supervision İ.Ş.S.; Resource - Ö.D., S.G. Materials - Ö.D., S.G.; Data Collection and/or Processing - Ö.D., S.G.; Analysis and/or Interpretation - İ.Ş.S., Ö.D.; Literature Search - İ.Ş.S, Ö.D; Writing - İ.Ş.S., Ö.D.; Critical Reviews - İ.Ş.S., Ö.D.

Conflict of Interest: Authors have no conflicts of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Bu çalışma için etik komite onayı Çukurova Üniversitesi Kurumsal İnceleme Kurulu'ndan alınmıştır (CUM 2015-48-28).

Hasta Onamı: Hasta onamı bu çalışmaya katılan hastalardan alınmıştır.

Hakem Değerlendirmesi: Dış Bağımsız.

Yazar Katkıları: Fikir - Ö.D., İ.Ş.S.; Tasarım - İ.Ş.S., Ö.D., S.G.; Denetleme - İ.Ş.S.; Kaynaklar - Ö.D., S.G.; Malzemeler - Ö.D., S.G.; Veri Toplanması ve/veya işleme - Ö.D., S.G.; Analiz ve/veya Yorum - İ.Ş.S., Ö.D.; Literatür taraması - İ.Ş.S., Ö.D.; Yazıyı Yazan - İ.Ş.S., Ö.D.; Eleştirel İnceleme - İ.Ş.S., Ö.D.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

References

1. Sugarbaker PH, Cunliffe WJ, Belliveau J, de Bruijn EA, Graves T, Mullins RE, et al. Rationale for integrating early postoperative intraperitoneal chemotherapy into the surgical treatment of gastrointestinal cancer. *Semin Oncol* 1989; 16: 83-97.
2. Chua TC, Yan TD, Saxena A, Morris DL. Should the treatment of peritoneal carcinomatosis by cytoreductive surgery and hyperthermic intraperitoneal chemotherapy still be regarded as a highly morbid procedure?: a systematic review of morbidity and mortality. *Ann Surg* 2009; 249: 900-7. [\[CrossRef\]](#)
3. Wilson KA, Dowling AJ, Abdolell M, Tannock IF. Perception of quality of life by patients, partners and treating physicians. *Qual Life Res* 2000; 9: 1041-52. [\[CrossRef\]](#)
4. Jacquet P, Sugarbaker PH. Current methodologies for clinical assessment of patients with peritoneal carcinomatosis. *J Exp Clin Cancer Res* 1996; 15: 49-58.
5. Esquivel J, Elias D, Baratti D, Kusamura S, Deraco M. Consensus statement on the loco regional treatment of colorectal cancer with peritoneal dissemination. *J Surg Oncol* 2008; 98: 263-7. [\[CrossRef\]](#)
6. Hoopman R, Muller MJ, Terwee CB, Aaronson NK. Translation and validation of the EORTC QLQ-C30 for use among Turkish and Moroccan ethnic minority cancer patients in the Netherlands. *Eur J Cancer* 2006; 42: 1839-47. [\[CrossRef\]](#)
7. Fayers PM, Aaronson NK, Bjordal K, Curran D, Groenvold M. The EORTC QLQ-C30 Scoring Manual. 2nd Ed., Brussels: In: European Organization for Research and Treatment of Cancer; 1999.
8. Schmidt U, Dahlke MH, Klempnauer J, Schlitt HJ, Piso P. Perioperative morbidity and quality of life in long-term survivors following cytoreductive surgery and hyperthermic intraperitoneal chemotherapy. *Eur J Surg Oncol*. 2005; 31: 53-8. [\[CrossRef\]](#)
9. Alves S, Mohamed F, Yadegarfar G, Youssef H, Moran BJ. Prospective longitudinal study of quality of life following cytoreductive surgery and intraperitoneal chemotherapy for pseudomyxoma peritonei. *Eur J Surg Oncol* 2010; 36: 1156-61. [\[CrossRef\]](#)
10. Tan WJ, Wong JF, Chia CS, Tan GH, Soo KC, Teo MC. Quality of life after cytoreductive surgery and hyperthermic intraperitoneal chemotherapy: an Asian perspective. *Ann Surg Oncol* 2013; 20: 4219-23. [\[CrossRef\]](#)

Cite this article as: Düzgün Ö, Sarıcı İŞ, Gökçay S. Short-term Quality of Life after Cytoreductive Surgery and Hyperthermic Intraperitoneal Chemotherapy. *Istanbul Med J* 2018; 19: 143-6.